

Clinical Policy: Omacetaxine (Synribo)

Reference Number: CP.PHAR.108

Effective Date: 04.01.13

Last Review Date: 05.19

Line of Business: Commercial, HIM, Medicaid

[Coding Implications](#)

[Revision Log](#)

See [Important Reminder](#) at the end of this policy for important regulatory and legal information.

Description

Omacetaxine (Synribo[®]) is cephalotaxine ester that inhibits protein synthesis by binding to the A-site in the peptidyl-transferase center of the large ribosomal subunit.

FDA Approved Indication(s)

Synribo is indicated for the treatment of adult patients with chronic or accelerated phase chronic myeloid leukemia (CML) with resistance and/or intolerance to two or more tyrosine kinase inhibitors.

Policy/Criteria

Provider must submit documentation (such as office chart notes, lab results or other clinical information) supporting that member has met all approval criteria.

It is the policy of health plans affiliated with Centene Corporation[®] that Synribo is **medically necessary** when the following criteria are met:

I. Initial Approval Criteria

A. Chronic Myeloid Leukemia (must meet all):

1. Diagnosis of Ph⁺ (BCR-ABL1-positive) CML;
2. Prescribed by or in consultation with an oncologist or hematologist;
3. Age \geq 18 years;
4. Request meets one of the following (a or b):
 - a. Dose does not exceed 2.5 mg/m² per day.
 - b. Dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (prescriber must submit supporting evidence).

Approval duration:

Medicaid/HIM - 6 months

Commercial - 6 months or to the member's renewal date, whichever is longer

B. Other diagnoses/indications

1. Refer to the off-label use policy for the relevant line of business if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized): CP.CPA.09 for commercial, HIM.PHAR.21 for health insurance marketplace, and CP.PMN.53 for Medicaid.

II. Continued Therapy

A. Chronic Myeloid Leukemia (must meet all):

1. Currently receiving medication via Centene benefit, or documentation supports that member is currently receiving Synribo for CML and has received this medication for at least 30 days;
2. Member is responding positively to therapy;
3. If request is for a dose increase, request meets one of the following (a or b):
 - a. New dose does not exceed 2.5 mg/m² per day;
 - b. New dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).

Approval duration:

Medicaid/HIM - 12 months

Commercial - 6 months or to the member's renewal date, whichever is longer

B. Other diagnoses/indications (must meet 1 or 2):

1. Currently receiving medication via Centene benefit and documentation supports positive response to therapy.

Approval duration: Duration of request or 6 months (whichever is less); or

2. Refer to the off-label use policy for the relevant line of business if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized): CP.CPA.09 for commercial, HIM.PHAR.21 for health insurance marketplace, and CP.PMN.53 for Medicaid.

III. Diagnoses/Indications for which coverage is NOT authorized:

- ### A. Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off label use policies – CP.CPA.09 for commercial, HIM.PHAR.21 for health insurance marketplace, and CP.PMN.53 for Medicaid or evidence of coverage documents.

IV. Appendices/General Information

Appendix A: Abbreviation/Acronym Key

CML: chronic myelogenous leukemia

FDA: Food and Drug Administration

Appendix B: Therapeutic Alternatives

Not applicable

Appendix C: Contraindications/Boxed Warnings

- Contraindication(s): none reported
- Boxed warning(s): arterial occlusion, venous thromboembolism, heart failure, hepatotoxicity.

V. Dosage and Administration

Indication	Dosing Regimen	Maximum Dose
CML	Induction dose: 1.25 mg/m ² subcutaneous twice daily for 14 consecutive days of a 28-day cycle	2.5 mg/m ² per day

Indication	Dosing Regimen	Maximum Dose
	Maintenance dose: 1.25 mg/m ² subcutaneous twice daily for 7 consecutive days of a 28-day cycle	

VI. Product Availability

Single-use vial: 3.5 mg of omacetaxine mepesuccinate as a lyophilized powder

VII. References

1. Synribo Prescribing Information. North Wales, PA: Teva Pharmaceuticals USA, Inc.; June 2017. Available at http://www.synribohcp.com/pdf/synribo_pi.pdf. Accessed February 5, 2019.
2. National Comprehensive Cancer Network Drugs and Biologics Compendium. Available at www.nccn.org. Accessed February 5, 2019.
3. National Comprehensive Cancer Network Guidelines. Chronic Myeloid Leukemia Version 1.2019. Available at www.nccn.org. Accessed February 5, 2019.

Coding Implications

Codes referenced in this clinical policy are for informational purposes only. Inclusion or exclusion of any codes does not guarantee coverage. Providers should reference the most up-to-date sources of professional coding guidance prior to the submission of claims for reimbursement of covered services.

HCPCS Codes	Description
J9262	Injection, omacetaxine mepesuccinate, 0.01 mg

Reviews, Revisions, and Approvals	Date	P&T Approval Date
Policy converted to new template. Criteria: initial approval period shortened to three months per NCCN monitoring recommendation starting three months post therapy change; documentation requests removed and replaced with attestation requests; denial based on myelosuppression removed; detailed efficacy criteria removed and replaced with general disease progression criteria.	11.15	12.15
Policy converted to new template. Removed prescriber and age requirements. Added NCCN recommended use (CML relapse post-transplantation). Removed attestation that member does not have poorly controlled diabetes from initial criteria.	11.16	12.16
Changed approval durations from 3/6 months to 6/12 months; Added age; Added NCCN recommended uses of previously diagnosed with chronic phase CML and has progressed to accelerated phase CML and history of T315I mutation.	07.01.17	11.17
2Q 2018 annual review: no significant changes; added Commercial and HIM lines of business; added continuity of care statement;	02.13.18	05.18

Reviews, Revisions, and Approvals	Date	P&T Approval Date
summarized NCCN and FDA approved uses for improved clarity; added specialist involvement in care; references reviewed and updated.		
2Q 2019 annual review: Ph+ designation added to CML; hematologist added to CML/ALL criteria; references reviewed and updated.	02.19.19	05.19

Important Reminder

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. The Health Plan makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved. “Health Plan” means a health plan that has adopted this clinical policy and that is operated or administered, in whole or in part, by Centene Management Company, LLC, or any of such health plan’s affiliates, as applicable.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable Health Plan-level administrative policies and procedures.

This clinical policy is effective as of the date determined by the Health Plan. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. The Health Plan retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members. This clinical policy is not intended to recommend treatment for members. Members should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom the Health Plan has no control or right of control. Providers are not agents or employees of the Health Plan.

This clinical policy is the property of the Health Plan. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members and their representatives agree to be bound by such terms and conditions by providing services to members and/or submitting claims for payment for such services.

Note:

For Medicaid members, when state Medicaid coverage provisions conflict with the coverage provisions in this clinical policy, state Medicaid coverage provisions take precedence. Please refer to the state Medicaid manual for any coverage provisions pertaining to this clinical policy.

For Health Insurance Marketplace members, when applicable, this policy applies only when the prescribed agent is on your health plan approved formulary. Request for non-formulary drugs must be reviewed using the formulary exception policy.

©2013 Centene Corporation. All rights reserved. All materials are exclusively owned by Centene Corporation and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Centene Corporation. You may not alter or remove any trademark, copyright or other notice contained herein. Centene[®] and Centene Corporation[®] are registered trademarks exclusively owned by Centene Corporation.