

Clinical Policy: Idelalisib (Zydelig)

Reference Number: CP.PHAR.133

Effective Date: 12.01.18

Last Review Date: 11.20

Line of Business: Commercial, HIM, Medicaid

[Revision Log](#)

See [Important Reminder](#) at the end of this policy for important regulatory and legal information.

Description

Idelalisib (Zydelig[®]) is a kinase inhibitor.

FDA Approved Indication(s)

Zydelig is indicated for the treatment of:

- Relapsed chronic lymphocytic leukemia (CLL), in combination with rituximab, in patients for whom rituximab alone would be considered appropriate therapy due to other co-morbidities
- Relapsed follicular B-cell non-Hodgkin lymphoma (FL) in patients who have received at least two prior systemic therapies*
- Relapsed small lymphocytic lymphoma (SLL) in patients who have received at least two prior systemic therapies*

**Accelerated approval was granted for FL and SLL based on overall response rate. Improvement in patient survival or disease related symptoms has not been established. Continued approval for these indications may be contingent upon verification of clinical benefit in confirmatory trials.*

Limitation(s) of use:

- Zydelig is not indicated and is not recommended for first-line treatment of any patient.
- Zydelig is not indicated and is not recommended in combination with bendamustine and/or rituximab for the treatment of FL.

Policy/Criteria

Provider must submit documentation (such as office chart notes, lab results or other clinical information) supporting that member has met all approval criteria.

It is the policy of health plans affiliated with Centene Corporation[®] that Zydelig is **medically necessary** when the following criteria are met:

I. Initial Approval Criteria

A. Chronic Lymphocytic Leukemia/Small Lymphocytic Lymphoma (must meet all):

1. Diagnosis of CLL or SLL;
2. Prescribed by or in consultation with an oncologist or hematologist;
3. Age \geq 18 years;
4. Relapsed/refractory disease after \geq one prior therapy (*see Appendix B for examples*);*
**Prior authorization may be required.*
5. Request meets one of the following (a or b):*
 - a. Dose does not exceed 300 mg per day (2 tablets per day);

- b. Dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).

**Prescribed regimen must be FDA-approved or recommended by NCCN.*

Approval duration:

Medicaid/HIM – 6 months

Commercial – Length of Benefit

B. Follicular and Marginal Zone Lymphomas (must meet all):

1. One of the following diagnoses (a or b):
 - a. FL;
 - b. Marginal zone lymphoma (off-label) (i, ii, or iii):
 - i. Splenic marginal zone lymphoma;
 - ii. Nodal marginal zone lymphoma;
 - iii. Extranodal marginal zone lymphoma (a or b):
 - a) Gastric MALT lymphoma;
 - b) Nongastric MALT lymphoma;
2. Prescribed by or in consultation with an oncologist or hematologist;
3. Age \geq 18 years;
4. Relapsed/refractory disease after \geq 2 prior therapies (*see Appendix B for examples*);*
**Prior authorization may be required.*
5. Request meets one of the following (a or b):*
 - a. Dose does not exceed 300 mg per day (2 tablets per day);
 - b. Dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).

**Prescribed regimen must be FDA-approved or recommended by NCCN.*

Approval duration:

Medicaid/HIM – 6 months

Commercial – Length of Benefit

C. Other diagnoses/indications

1. Refer to the off-label use policy for the relevant line of business if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized): CP.CPA.09 for commercial, HIM.PHAR.21 for health insurance marketplace, and CP.PMN.53 for Medicaid.

II. Continued Therapy

A. All Indications in Section I (must meet all):

1. Currently receiving medication via Centene benefit, or documentation supports that member is currently receiving Zydelig for a covered indication and has received this medication for at least 30 days;
2. Member is responding positively to therapy;
3. If request is for a dose increase, request meets one of the following (a or b):*
 - a. New dose does not exceed 300 mg per day (2 tablets per day);
 - b. New dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).

**Prescribed regimen must be FDA-approved or recommended by NCCN.*

Approval duration:

Medicaid/HIM – 12 months
Commercial – Length of Benefit

B. Other diagnoses/indications (must meet 1 or 2):

1. Currently receiving medication via Centene benefit and documentation supports positive response to therapy.
Approval duration: Duration of request or 6 months (whichever is less); or
2. Refer to the off-label use policy for the relevant line of business if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized): CP.CPA.09 for commercial, HIM.PHAR.21 for health insurance marketplace, and CP.PMN.53 for Medicaid.

III. Diagnoses/Indications for which coverage is NOT authorized:

- A.** Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off label use policies – CP.CPA.09 for commercial, HIM.PHAR.21 for health insurance marketplace, and CP.PMN.53 for Medicaid or evidence of coverage documents.

IV. Appendices/General Information

Appendix A: Abbreviation/Acronym Key

CLL: chronic lymphocytic leukemia	NCCN: National Comprehensive Cancer Network
FDA: Food and Drug Administration	SLL: small lymphocytic lymphoma
FL: follicular B-cell non-Hodgkin lymphoma	

Appendix B: Therapeutic Alternatives

This table provides a listing of preferred alternative therapy recommended in the approval criteria. The drugs listed here may not be a formulary agent for all relevant lines of business and may require prior authorization.

Drug Name	Dosing Regimen	Dose Limit/ Maximum Dose
<p>CLL/SLL <i>Examples of first-line, second-line and subsequent therapies:</i></p> <ul style="list-style-type: none"> • FCR (fludarabine, cyclophosphamide, rituximab) • HDMP (high-dose methylprednisolone) + rituximab • Single-agent examples: Imbruvica[®] (ibrutinib); Venclexta[®] (venetoclax) ± Gazyva[®] (obinutuzumab) or rituximab; Campath[®] (alemtuzumab) ± rituximab; Gazyva; Copiktra[®] (duvelisib); Calquence[®] (acalabrutinib); Revlimid[®] (lenalidomide) ± rituximab; Arzerra[®] (ofatumumab) ± FC (fludarabine, cyclophosphamide); Leukeran[®] (chlorambucil) + rituximab 	Varies	Varies
<p>Follicular Lymphoma <i>Examples of first-line, second-line and subsequent therapies:</i></p> <ul style="list-style-type: none"> • bendamustine + Gazyva or rituximab 	Varies	Varies

Drug Name	Dosing Regimen	Dose Limit/ Maximum Dose
<ul style="list-style-type: none"> CHOP (cyclophosphamide, doxorubicin, vincristine, predenisonone) + Gazyva or rituximab CVP (cyclophosphamide, vincristine, prednisone) + Gazyva or rituximab <u>Single-agent examples</u>: rituximab; Revlimid ± rituximab 		
<p>Marginal Zone Lymphomas <i>Examples of first-line, second-line and subsequent therapies:</i></p> <ul style="list-style-type: none"> bendamustine + rituximab RCHOP (rituximab, cyclophosphamide, doxorubicin, vincristine, prednisone) RCVP (rituximab, cyclophosphamide, vincristine, prednisone) <u>Single-agent examples</u>: rituximab; Leukeran ± rituximab; cyclophosphamide ± rituximab; Imbruvica; Revlimid ± rituximab; Copiktra; Aliqopa[®] (copanlisib) 	Varies	Varies

Therapeutic alternatives are listed as Brand name[®] (generic) when the drug is available by brand name only and generic (Brand name[®]) when the drug is available by both brand and generic.

Appendix C: Contraindications/Boxed Warnings

- Contraindication(s): history of serious allergic reactions including anaphylaxis and toxic epidermal necrolysis
- Boxed warning(s): fatal and serious toxicities - hepatic, severe diarrhea, colitis, pneumonitis, infections, and intestinal perforation

V. Dosage and Administration

Indication	Dosing Regimen	Maximum Dose
CLL, FL, SLL	150 mg PO BID	300 mg per day

VI. Product Availability

Tablets: 150 mg, 100 mg

VII. References

- Zydelig Prescribing Information. Foster City, CA: Gilead Sciences, Inc.; October 2018. Available at <http://www.zydelig.com>. Accessed July 24, 2020.
- National Comprehensive Cancer Network Drugs and Biologics Compendium. Available at: http://www.nccn.org/professionals/drug_compendium. Accessed July 24, 2020.
- National Comprehensive Cancer Network. Chronic Lymphocytic Leukemia/Small Lymphocytic Lymphoma Version 4.2020. Available at: https://www.nccn.org/professionals/physician_gls/pdf/ll.pdf. Accessed July 24, 2020.
- National Comprehensive Cancer Network. B-cell lymphomas Version 2.2020. Available at www.nccn.org. Accessed July 24, 2020.

Reviews, Revisions, and Approvals	Date	P&T Approval Date
Policy created: adapted from previously approved HIM.PA.SP49 and CP.CPA.278 Idelalisib (Zydelig) (both to be retired); new policy for Centene Medicaid; summarized NCCN and FDA-approved uses for improved clarity; added age requirement and specialist involvement in care; removed primary cutaneous B-cell lymphoma as a covered off-label indication (disease not listed in the NCCN compendium for Zydelig); updated continued therapy section to include language for continuity of care; references reviewed and updated.	07.11.18	11.18
4Q 2019 annual review: Criteria/Appendix B reorganized to reconcile with similar policies; FDA/NCCN dosing limitation added, references reviewed and updated.	08.27.19	11.19
4Q 2020 annual review: no significant changes; references reviewed and updated.	08.11.20	11.20

Important Reminder

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. The Health Plan makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved. “Health Plan” means a health plan that has adopted this clinical policy and that is operated or administered, in whole or in part, by Centene Management Company, LLC, or any of such health plan’s affiliates, as applicable.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable Health Plan-level administrative policies and procedures.

This clinical policy is effective as of the date determined by the Health Plan. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. The Health Plan retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members. This clinical policy is not intended to recommend treatment for members. Members should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom the Health Plan has no control or right of control. Providers are not agents or employees of the Health Plan.

This clinical policy is the property of the Health Plan. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members and their representatives agree to be bound by such terms and conditions by providing services to members and/or submitting claims for payment for such services.

Note:

For Medicaid members, when state Medicaid coverage provisions conflict with the coverage provisions in this clinical policy, state Medicaid coverage provisions take precedence. Please refer to the state Medicaid manual for any coverage provisions pertaining to this clinical policy.

©2018 Centene Corporation. All rights reserved. All materials are exclusively owned by Centene Corporation and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Centene Corporation. You may not alter or remove any trademark, copyright or other notice contained herein. Centene[®] and Centene Corporation[®] are registered trademarks exclusively owned by Centene Corporation.