

## Clinical Policy: Methotrexate (Otrexup, Rasuvo, Xatmep, Reditrex)

Reference Number: CP.PHAR.134

Effective Date: 12.01.18

Last Review Date: 11.20

<u>Coding Implications</u>

<u>Revision Log</u>

Line of Business: Commercial, HIM, Medicaid

See <u>Important Reminder</u> at the end of this policy for important regulatory and legal information.

## **Description**

Methotrexate injection (Otrexup<sup>TM</sup>, Rasuvo<sup>®</sup>, Reditrex<sup>TM</sup>) and oral solution (Xatmep<sup>®</sup>) are folate analog metabolic inhibitors.

## FDA Approved Indication(s)

Otrexup, Rasuvo, and Reditrex are indicated for:

- Management of selected adults with severe, active rheumatoid arthritis (RA), or children with active polyarticular juvenile idiopathic arthritis (pJIA), who have had an insufficient therapeutic response to, or are intolerant of, an adequate trial of first-line therapy including full dose non-steroidal anti-inflammatory agents (NSAIDs)
- In adults for the symptomatic control of severe, recalcitrant, disabling psoriasis (PsO) that is not adequately responsive to other forms of therapy, but only when the diagnosis has been established, as by biopsy and/or after dermatologic consultation

Limitation(s) of use: Otrexup, Rasuvo, and Reditrex are not indicated for the treatment of neoplastic diseases.

### Xatmep is indicated for:

- Treatment of pediatric patients with acute lymphoblastic leukemia (ALL) as part of a multiphase, combination chemotherapy maintenance regimen
- Management of pediatric patients with active polyarticular juvenile idiopathic arthritis (pJIA)
  who have had an insufficient therepeutic response to, or are intolerant of, an adequate trial of
  first-line therapy including full dose NSAIDs

### Policy/Criteria

Provider must submit documentation (such as office chart notes, lab results or other clinical information) supporting that member has met all approval criteria.

It is the policy of health plans affiliated with Centene Corporation<sup>®</sup> that Otrexup, Rasuvo, Xatmep, and Reditrex are **medically necessary** when the following criteria are met:

#### I. Initial Approval Criteria

- A. Polyarticular Juvenile Idiopathic Arthritis (must meet all):
  - 1. Diagnosis of PJIA;
  - 2. Prescribed by or in consultation with a rheumatologist;
  - 3. Member meets one of the following (a or b):
    - a. For Otrexup, Rasuvo, or Reditrex: age  $\geq 2$  years;



- b. For Xatmep: age  $\leq 18$  years;
- 4. For Otrexup, Rasuvo, or Reditrex: failure of generic methotrexate injection, unless contraindicated or clinically significant adverse effects are experienced;
- 5. For Xatmep: documentation supports inability to swallow pills;
- 6. Dose does not exceed the following (a or b):
  - a. Otrexup, Ravuso, or Reditrex: 20 mg per week;
  - b. Xatmep: 30 mg/m<sup>2</sup> per week.

### **Medicaid/HIM** – 6 months

Commercial – 6 months or to the member's renewal date, whichever is longer

#### B. Rheumatoid Arthritis or Psoriasis (must meet all):

- 1. Diagnosis of RA or PsO;
- 2. Request is for Otrexup, Rasuvo, or Reditrex;
- 3. For RA: prescribed by or in consultation with a rheumatologist;
- 4. For PsO: by or in consultation with a rheumatologist or a dermatologist;
- 5. Age  $\geq 2$  years;
- 6. Failure of generic methotrexate injection, unless contraindicated or clinically significant adverse effects are experienced;
- 7. Dose does not exceed the following (a or b):
  - a. RA: 20 mg per week;
  - b. Psoriasis: 30 mg per week.

### **Medicaid/HIM** – 6 months

Commercial – 6 months or to the member's renewal date, whichever is longer

## C. Acute Lymphoblastic Leukemia (must meet all):

- 1. Diagnosis of ALL;
- 2. Request is for Xatmep;
- 3. Prescribed by by or in consultation with an oncologist or hematologist;
- 4. Age < 18 years;
- 5. Documentation supports inability to swallow pills;
- 6. Request meets one of the following (a or b):\*
  - a. Dose does not exceed 30 mg/m<sup>2</sup> per week;
  - b. Dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).

\*Prescribed regimen must be FDA-approved or recommended by NCCN

### **Approval duration:**

**Medicaid/HIM** – 6 months

Commercial – Length of Benefit

#### D. Other diagnoses/indications

1. Refer to the off-label use policy for the relevant line of business if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized): CP.CPA.09 for commercial, HIM.PHAR.21 for health insurance marketplace, and CP.PMN.53 for Medicaid.

## **II. Continued Therapy**

A. All Indications in Section I (must meet all):



- 1. Member meets one of the following (a or b):
  - a. Currently receiving medication via Centene benefit or member has previously met initial approval criteria;
  - b. Documentation supports that member is currently receiving Xatmep for ALL and has received this medication for at least 30 days
- 2. Member is responding positively to therapy;
- 3. If request is for a dose increase, new dose does not exceed the following (a or b):
  - a. Otrexup, Ravuso, or Reditrex:
    - i. RA, pJIA: 20 mg per week;
    - ii. Psoriasis: 30 mg per week;
  - b. Xatmep:
    - i. pJIA: 30 mg/m<sup>2</sup> per week;
    - ii. ALL: Request meets one of the following (1 or 2):\*
      - 1. Dose does not exceed 20 mg/m<sup>2</sup> per week;
      - 2. Dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).

\*Prescribed regimen must be FDA-approved or recommended by NCCN.

## Approval duration:

**Medicaid/HIM** – 12 months

**Commercial** – Otrexup, Rasuvo, and Reditrex: 6 months or to the member's renewal date, whichever is longer;

Xatmep: Length of Benefit

## **B.** Other diagnoses/indications (must meet 1 or 2):

1. Currently receiving medication via Centene benefit and documentation supports positive response to therapy.

## Approval duration: Duration of request or 12 months (whichever is less); or

2. Refer to the off-label use policy for the relevant line of business if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized): CP.CPA.09 for commercial, HIM.PHAR.21 for health insurance marketplace, and CP.PMN.53 for Medicaid.

#### III. Diagnoses/Indications for which coverage is NOT authorized:

**A.** Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off-label use policy – CP.CPA.09 for commercial, HIM.PHAR.21 for health insurance marketplace, and CP.PMN.53 for Medicaid, or evidence of coverage documents.

#### IV. Appendices/General Information

Appendix A: Abbreviation Key
ALL: acute lymphoblastic leukemia
FDA: Food and Drug Administration

NSAID: non-steroidal anti-inflammatory drug

PJIA: polyarticular juvenile idiopathic arthritis

PsO: psoriasis

RA: rheumatoid arthritis



Appendix B: Therapeutic Alternatives

Drug Name	Dosing Regimen	Dose Limit/ Maximum Dose
methotrexate injection	RA 7.5 mg SC once weekly	RA, pJIA: 20 mg/week; PsO: 30 mg/week
	PJIA 10 mg/m <sup>2</sup> SC once weekly	1 30. 30 mg/week
	PsO 10-25 mg SC once weekly	
methotrexate tablets	ALL, PJIA 10 – 30 mg/m <sup>2</sup> once weekly	30 mg/m²/week

Therapeutic alternatives are listed as Brand name<sup>®</sup> (generic) when the drug is available by brand name only and generic (Brand name<sup>®</sup>) when the drug is available by both brand and generic.

## Appendix C: Contraindications/Boxed Warnings

- Contraindication(s):
  - o Otrexup, Rasuvo, Reditrex: pregnancy; alcoholism or liver disease; immunodeficiency syndromes; pre-existing blood dyscrasias; hypersensitivity
  - Xatmep: pregnancy in patients with non-malignant diseases; severe hypersensitivity to methotrexate
- Boxed warning(s): severe toxic reactions, including embryo-fetal toxicity and death

### Appendix D: General Information

• Otrexup, Rasuvo, and Reditrex are not indicated for the treatment of neoplastic diseases.

## V. Dosage and Administration

Drug Name	Indication	Dosing Regimen	Maximum Dose
Methotrexate	RA	7.5 mg SC once weekly	20 mg/week
injection	PJIA	10 mg/m <sup>2</sup> SC once weekly	20 mg/week
(Otrexup,			
Rasuvo,	PsO	10-25 mg SC once weekly	30 mg/week
Reditrex)			
Methotrexate	ALL	20 mg/m <sup>2</sup> PO once weekly	20 mg/m <sup>2</sup> /week
oral solution (Xatmep)	PJIA	10 mg/m <sup>2</sup> PO once weekly	30 mg/m <sup>2</sup> /week

### VI. Product Availability

Drug	Availability
Methotrexate injection	Auto-injector: 10 mg/0.4 mL, 12.5 mg/0.4 mL, 15 mg/0.4 mL, 17.5
(Otrexup)	mg/0.4 mL, 20 mg/0.4 mL, 22.5 mg/0.4 mL, 25 mg/0.4 mL
Methotrexate	Auto-injector: 7.5 mg/0.15 mL, 10 mg/0.2 mL, 12.5 mg/0.25 mL, 15
injection	mg/0.3 mL, 17.5 mg/0.35 mL, 20 mg/0.4 mL, 22.5 mg/0.45 mL, 25
(Rasuvo)	mg/0.5 mL, 27.5 mg/0.55 mL, 30 mg/0.6 mL



Drug	Availability
Methotrexate	Single-dose pre-filled injection: 7.5 mg/0.3 mL, 10 mg/0.4 mL, 12.5
injection	mg/0.5 mL, 15 mg/0.6 mL, 17.5 mg/0.7 mL, 20 mg/0.8 mL, 22.5
(Reditrex)	mg/0.9 mL, 25 mg/mL
Methotrexate oral	
solution	2.5 mg/mL in a 60 mL or 120 mL bottle
(Xatmep)	

#### VII. References

- 1. Otrexup Prescribing Information. Ewing, NJ: Antares Pharma, Inc. December 2019. Available at: <a href="https://www.otrexup.com">www.otrexup.com</a>. Accessed August 3, 2020.
- 2. Rasuvo Prescribing Information. Chicago, IL: Medac Pharma, Inc. March 2020. Availale at: <a href="http://cdn.rasuvo.com/assets/pdf/Prescribing-Information-current.pdf">http://cdn.rasuvo.com/assets/pdf/Prescribing-Information-current.pdf</a>. Accessed August 3, 2020.
- 3. Xatmep Prescribing Information. Greenwood Village, CO: Silvergate Pharmaceuticals, Inc.; March 2020. Available at: <a href="https://www.xatmep.com">www.xatmep.com</a>. Accessed August 3, 2020.
- Reditrex Prescribing Information. Nashville, TN: Cumberland Pharmaceuticals, Inc.; November 2019. Available at: <a href="https://www.accessdata.fda.gov/drugsatfda\_docs/label/2019/210737s000lbl.pdf">https://www.accessdata.fda.gov/drugsatfda\_docs/label/2019/210737s000lbl.pdf</a>. Accessed August 3, 2020.
- 5. Clinical Pharmacology [database online]. Tampa, FL: Gold Standard, Inc.; 2018. Available at: http://www.clinicalpharmacology-ip.com/. Accessed August 3, 2020.

## **Coding Implications**

Codes referenced in this clinical policy are for informational purposes only. Inclusion or exclusion of any codes does not guarantee coverage. Providers should reference the most up-to-date sources of professional coding guidance prior to the submission of claims for reimbursement of covered services.

HCPCS	Description
Codes	
J9250	Methotrexate sodium, 5 mg
J9260	Methotrexate sodium, 50 mg

Reviews, Revisions, and Approvals		P&T
		Approval
		Date
Policy created: adapted from CP.CPA.193 added Medicaid and HIM	07.31.18	11.18
Medical added; added Xatmep to policy; added specialist requirement;		
added age limits; references reviewed and updated.		
No significant changes; revised off-label use policy references to	06.24.19	
relevant lines of business.		
4Q 2019 annual review: no significant changes; added HIM NF		11.19
language for Xatmep; references reviewed and updated.		
RT4: added Reditrex to policy; revised HIM-medical benefit to HIM	03.17.20	
line of business.		



Reviews, Revisions, and Approvals		P&T
		Approval Date
4Q 2020 annual review: no significant changes; references reviewed and updated.	08.03.20	11.20

#### **Important Reminder**

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. The Health Plan makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved. "Health Plan" means a health plan that has adopted this clinical policy and that is operated or administered, in whole or in part, by Centene Management Company, LLC, or any of such health plan's affiliates, as applicable.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable Health Plan-level administrative policies and procedures.

This clinical policy is effective as of the date determined by the Health Plan. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. The Health Plan retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members. This clinical policy is not intended to recommend treatment for members. Members should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom the Health Plan has no control or right of control. Providers are not agents or employees of the Health Plan.



This clinical policy is the property of the Health Plan. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members and their representatives agree to be bound by such terms and conditions by providing services to members and/or submitting claims for payment for such services.

#### Note:

**For Medicaid members**, when state Medicaid coverage provisions conflict with the coverage provisions in this clinical policy, state Medicaid coverage provisions take precedence. Please refer to the state Medicaid manual for any coverage provisions pertaining to this clinical policy.

©2016 Centene Corporation. All rights reserved. All materials are exclusively owned by Centene Corporation and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Centene Corporation. You may not alter or remove any trademark, copyright or other notice contained herein. Centene® and Centene Corporation® are registered trademarks exclusively owned by Centene Corporation.