

Clinical Policy: Enasidenib (Idhifa)

Reference Number: CP.PHAR.363

Effective Date: 09.05.17 Last Review Date: 11.20

Line of Business: Commercial, HIM, Medicaid

Revision Log

See <u>Important Reminder</u> at the end of this policy for important regulatory and legal information.

Description

Enasidenib (Idhifa®) is an isocitrate dehydrogenase-2 (IDH2) inhibitor.

FDA Approved Indication(s)

Idhifa is indicated for the treatment of adult patients with relapsed or refractory acute myeloid leukemia (AML) with an IDH2 mutation as detected by an FDA-approved test.

Policy/Criteria

Provider must submit documentation (such as office chart notes, lab results or other clinical information) supporting that member has met all approval criteria.

It is the policy of health plans affiliated with Centene Corporation[®] that Idhifa is **medically necessary** when the following criteria are met:

I. Initial Approval Criteria

- A. Acute Myeloid Leukemia (must meet all):
 - 1. Diagnosis of AML;
 - 2. Prescribed by or in consultation with an oncologist or hematologist;
 - 3. Age \geq 18 years;
 - 4. Member meets one of the following (a, b, or c):
 - a. Age \geq 60 years;
 - b. Disease has relapsed after or is in remission following Idhifa therapy;
 - c. Disease has relapsed after or is refractory to induction therapy (see Appendix B for examples);*
 - *Prior authorization may be required.
 - 5. Presence of an IDH2 mutation;
 - 6. Request meets one of the following (a or b):*
 - a. Dose does not exceed 100 mg (1 tablet) per day;
 - b. Dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).

*Prescribed regimen must be FDA-approved or recommended by NCCN.

Approval duration:

Medicaid/HIM – 6 months

Commercial – Length of Benefit

CLINICAL POLICY Enasidenib



B. Other diagnoses/indications

1. Refer to the off-label use policy for the relevant line of business if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized): CP.CPA.09 for commercial, HIM.PHAR.21 for health insurance marketplace and CP.PMN.53 for Medicaid.

II. Continued Therapy

A. Acute Myeloid Leukemia (must meet all):

- 1. Currently receiving medication via Centene benefit, or documentation supports that member is currently receiving Idhifa for AML and has received this medication for at least 30 days;
- 2. Member is responding positively to therapy;
- 3. If request is for a dose increase, request meets one of the following (a or b):*
 - a. New dose does not exceed 100 mg (1 tablet) per day;
 - b. New dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).

*Prescribed regimen must be FDA-approved or recommended by NCCN.

Approval duration:

Medicaid/HIM – 12 months

Commercial – Length of Benefit

B. Other diagnoses/indications(must meet 1 or 2):

- 1. Currently receiving medication via Centene benefit and documentation supports positive response to therapy.
 - Approval duration: Duration of request or 6 months (whichever is less); or
- 2. Refer to the off-label use policy for the relevant line of business if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized): CP.CPA.09 for commercial, HIM.PHAR.21 for health insurance marketplace and CP.PMN.53 for Medicaid.

III. Diagnoses/Indications for which coverage is NOT authorized:

A. Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off label use policies – CP.CPA.09 for commercial, HIM.PHAR.21 for health insurance marketplace and CP.PMN.53 for Medicaid or evidence of coverage documents.

IV. Appendices/General Information

Appendix A: Abbreviation/Acronym Key

AML: acute myeloid leukemia NCCN: National Comprehensive Cancer

FDA: Food and Drug Administration Network

IDH2: isocitrate dehydrogenase-2

CLINICAL POLICY Enasidenib



Appendix B: Therapeutic Alternatives

This table provides a listing of preferred alternative therapy recommended in the approval criteria. The drugs listed here may not be a formulary agent for all relevant lines of business

and may require prior authorization.

Drug Name	Dosing Regimen	Dose Limit/ Maximum Dose
cytarabine with idarubicin or daunorubicin	Age < 60 years: example of intensive induction therapy: cytarabine 100 – 200 mg/m² continuous IV infusion x 7 days with idarubicin 12 mg/m² IV or daunorubicin 60-90 mg/m² IV x 3 days	Varies
cytarabine with idarubicin or daunorubicin or mitoxantrone	Age \geq 60 years: example of intensive induction therapy: cytarabine $100-200 \text{ mg/m}^2$ continuous IV infusion x 7 days with idarubicin 12 mg/m² IV or daunorubicin 60-90 mg/m² IV x 3 days or mitoxantrone 12 mg/m² x 3 days	Varies

Therapeutic alternatives are listed as Brand name[®] (generic) when the drug is available by brand name only and generic (Brand name[®]) when the drug is available by both brand and generic.

Appendix C: Contraindications/Boxed Warnings

- Contraindication(s): none reported
- Boxed warning(s): differentiation syndrome. If differentiation syndrome is suspected, initiate corticosteroid therapy and hemodynamic monitoring until symptom resolution.

V. Dosage and Administration

Indication	Dosing Regimen	Maximum Dose
AML	100 mg PO QD	100 mg/day

VI. Product Availability

Tablets: 50 mg, 100 mg

VII. References

- 1. Idhifa Prescribing Information. Summit, NJ: Celgene Corporation; August 2019. Available at: www.idhifa.com. Accessed August 4, 2020.
- 2. National Comprehensive Cancer Network Drugs and Biologics Compendium. Available at: http://www.nccn.org/professionals/drug compendium. Accessed August 4, 2020.
- 3. National Comprehensive Cancer Network. Acute Myeloid Leukemia Version 3.2020. Available at: https://www.nccn.org/professionals/physician_gls/pdf/aml.pdf. Accessed August 4, 2020.

Reviews, Revisions, and Approvals	Date	P&T Approval Date
Policy created.	09.05.17	11.17

CLINICAL POLICY Enasidenib



Reviews, Revisions, and Approvals		P&T
		Approval
		Date
Combined Commercial and Medicaid policies.	02.12.18	
4Q 2018 annual review: specialist requirement was added; added	08.21.18	11.18
NCCN Compendium supported use in patients age ≥ 60 years who		
are not candidates for intensive remission induction therapy or		
declines intensive therapy; references reviewed and updated.		
4Q 2019 annual review: NCCN use added - relapse/remission post	08.27.19	11.19
Idhifa therapy; FDA/NCCN dosing limitation added; references		
reviewed and updated.		
Added HIM line of business.	02.13.20	
4Q 2020 annual review: no significant changes; updated Appendix C;	08.04.20	11.20
references reviewed and updated.		

Important Reminder

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. The Health Plan makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved. "Health Plan" means a health plan that has adopted this clinical policy and that is operated or administered, in whole or in part, by Centene Management Company, LLC, or any of such health plan's affiliates, as applicable.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable Health Plan-level administrative policies and procedures.

This clinical policy is effective as of the date determined by the Health Plan. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. The Health Plan retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise

CLINICAL POLICY Enasidenib



professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members. This clinical policy is not intended to recommend treatment for members. Members should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom the Health Plan has no control or right of control. Providers are not agents or employees of the Health Plan.

This clinical policy is the property of the Health Plan. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members and their representatives agree to be bound by such terms and conditions by providing services to members and/or submitting claims for payment for such services.

Note:

For Medicaid members, when state Medicaid coverage provisions conflict with the coverage provisions in this clinical policy, state Medicaid coverage provisions take precedence. Please refer to the state Medicaid manual for any coverage provisions pertaining to this clinical policy.

©2017 Centene Corporation. All rights reserved. All materials are exclusively owned by Centene Corporation and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Centene Corporation. You may not alter or remove any trademark, copyright or other notice contained herein. Centene® and Centene Corporation. The composition of the contained herein composition.