

**Clinical Policy: Rifamycin (Aemcolo)** 

Reference Number: CP.PMN.196

Effective Date: 06.01.19 Last Review Date: 05.21

Line of Business: Commercial, HIM, Medicaid Revision Log

See <u>Important Reminder</u> at the end of this policy for important regulatory and legal information.

# **Description**

Rifamycin (Aemcolo™) is an oral rifamycin antibacterial.

# FDA Approved Indication(s)

Aemcolo is indicated for the treatment of travelers' diarrhea (TD) caused by noninvasive strains of *Escherichia coli* in adults.

Limitation(s) of use: Aemcolo is not indicated in patients with diarrhea complicated by fever or bloody stool or due to pathogens other than noninvasive strains of *Escherichia coli*.

To reduce the development of drug-resistant bacteria and maintain the effectiveness of Aemcolo and other antibacterial drugs, Aemcolo should be used only to treat or prevent infections that are proven or strongly suspected to be caused by bacteria.

# Policy/Criteria

Provider must submit documentation (such as office chart notes, lab results or other clinical information) supporting that member has met all approval criteria.

It is the policy of health plans affiliated with Centene Corporation<sup>®</sup> that Aemcolo is **medically necessary** when the following criteria are met:

#### I. Initial Approval Criteria

- A. Travelers' Diarrhea (must meet all):
  - 1. Diagnosis of TD;
  - 2. Age  $\geq$  18 years;
  - 3. Failure of azithromycin 1,000 mg as a single dose, unless contraindicated or clinically significant adverse effects are experienced;
  - 4. Dose does not exceed 776 mg (4 tablets) per day.

Approval duration: 3 days

#### B. Other diagnoses/indications

1. Refer to the off-label use policy for the relevant line of business if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized): CP.CPA.09 for commercial, HIM.PA.154 for health insurance marketplace and CP.PMN.53 for Medicaid.

# CLINICAL POLICY Rifamycin



# **II. Continued Therapy**

#### A. Travelers' Diarrhea

1. May not be renewed as maximum allowed treatment duration is 3 days. Review initial approval criteria for new cases of travelers' diarrhea unrelated to original medication request.

Approval duration: Not applicable

### **B.** Other diagnoses/indications (must meet 1 or 2):

1. Currently receiving medication via Centene benefit and documentation supports positive response to therapy.

# Approval duration: Duration of request or 1 month (whichever is less); or

2. Refer to the off-label use policy for the relevant line of business if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized): CP.CPA.09 for commercial, HIM.PA.154 for health insurance marketplace and CP.PMN.53 for Medicaid.

### III. Diagnoses/Indications for which coverage is NOT authorized:

**A.** Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off label use policies – CP.CPA.09 for commercial, HIM.PA.154 for health insurance marketplace and CP.PMN.53 for Medicaid or evidence of coverage documents.

#### IV. Appendices/General Information

Appendix A: Abbreviation/Acronym Key FDA: Food and Drug Administration

TD: travelers' diarrhea

#### Appendix B: Therapeutic Alternatives

This table provides a listing of preferred alternative therapy recommended in the approval criteria. The drugs listed here may not be a formulary agent for all relevant lines of business and may require prior authorization.

Drug Name	Dosing Regimen	Dose Limit/ Maximum Dose
azithromycin	1,000 mg PO single	500 mg/day PO is FDA-approved dosage;
(Zithromax®)	dose	however, doses up to 1,200 mg/day PO are used
		off-label; 2 g PO when given as single dose

Therapeutic alternatives are listed as Brand name<sup>®</sup> (generic) when the drug is available by brand name only and generic (Brand name<sup>®</sup>) when the drug is available by both brand and generic.

#### Appendix C: Contraindications/Boxed Warnings

- Contraindication(s): known hypersensitivity to rifamycin, any of the other rifamycin class antimicrobial agents (e.g., rifaximin), or any of the components in Aemcolo
- Boxed warning(s): none reported

# CLINICAL POLICY Rifamycin



V. Dosage and Administration

Indication	<b>Dosing Regimen</b>	Maximum Dose
TD	388 mg PO BID for 3 days	776 mg/day

#### VI. Product Availability

Delayed-release tablet: 194 mg

#### VII. References

- 1. Aemcolo Prescribing Information. San Diego, CA: Aries Pharmaceuticals, Inc.; November 2018. Available at:
  - https://www.accessdata.fda.gov/drugsatfda\_docs/label/2018/210910s000lbl.pdf. Accessed February 17, 2021.
- 2. Connor BA. Centers for Disease Control and Prevention: Travelers' diarrhea, chapter 2 the pretravel consultation. Available at:
  - https://wwwnc.cdc.gov/travel/yellowbook/2020/preparing-international-travelers/travelers-diarrhea. Accessed February 17, 2021.
- 3. Riddle MS, et al. Guidelines for the prevention and treatment of travelers' diarrhea: a graded expert panel report. J Travel Med. 2017:24(Suppl 1):S63-80.
- 4. DuPont HL, et al. Targeting of rifamycin SV to the colon for treatment of travelers' diarrhea: a randomized, double-blind, placebo-controlled phase 3 study. J Travel Med. 2014:21(6):369–76.
- 5. Steffen R, Jiang Z, Garcia MLG, et al., Rifamycin SV-MMX for treatment of travelers' diarrhea: equally effective as ciprofloxacin and not associated with the acquisition of multi-drug resistant bacteria. J Travel Med. Tay116, <a href="https://doi.org/10.1093/jtm/tay116">https://doi.org/10.1093/jtm/tay116</a>. Published 20 November 2018.

Reviews, Revisions, and Approvals	Date	P&T Approval Date
Policy created	01.08.19	05.19
Requirement for a prior trial of a fluoroquinolone is removed due to concerns regarding increasing resistance to fluoroquinolones along with adverse dysbiotic (reduction in diversity of intestinal microbiota) and musculoskeletal adverse effects.	04.23.19	
Removed TBD HIM from included lines of business per SDC.	12.02.19	
2Q 2020 annual review: no significant changes; added HIM line of business; updated limitations of use; references reviewed and updated.	02.06.20	05.20
2Q 2021 annual review: no significant changes; revised reference to HIM off-label use policy from HIM.PHAR.21 to HIM.PA.154; references reviewed and updated.	02.17.21	05.21

# **Important Reminder**

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program

# CLINICAL POLICY Rifamycin



approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. The Health Plan makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved. "Health Plan" means a health plan that has adopted this clinical policy and that is operated or administered, in whole or in part, by Centene Management Company, LLC, or any of such health plan's affiliates, as applicable.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable Health Plan-level administrative policies and procedures.

This clinical policy is effective as of the date determined by the Health Plan. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. The Health Plan retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members. This clinical policy is not intended to recommend treatment for members. Members should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom the Health Plan has no control or right of control. Providers are not agents or employees of the Health Plan.

This clinical policy is the property of the Health Plan. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members and their representatives agree to be bound by such terms and conditions by providing services to members and/or submitting claims for payment for such services.

# CLINICAL POLICY Rifamycin



#### Note:

**For Medicaid members**, when state Medicaid coverage provisions conflict with the coverage provisions in this clinical policy, state Medicaid coverage provisions take precedence. Please refer to the state Medicaid manual for any coverage provisions pertaining to this clinical policy.

©2019 Centene Corporation. All rights reserved. All materials are exclusively owned by Centene Corporation and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Centene Corporation. You may not alter or remove any trademark, copyright or other notice contained herein. Centene® and Centene Corporation® are registered trademarks exclusively owned by Centene Corporation.